

# OBSERVATIONS & RECOMMENDATIONS

After reviewing data collected from **NUTTS POND, MANCHESTER**, the program coordinators have made the following observations and recommendations.

Thank you for your continued hard work sampling the pond this year! Your monitoring group sampled the deep spot **three** times this year and has done so for many years! As you know, conducting multiple sampling events each year enables DES to more accurately detect water quality changes. Keep up the good work!

In 2006, the City of Manchester received a DES Local Watershed Initiative Grant to improve the water quality of Nutts Pond by implementing stormwater Best Management Practices (BMPs) and Low Impact Development (LID) techniques. The proposed project will reduce the amount of non-point source pollutant loading to Nutts Pond from the eastern and southern sub-watersheds using structural BMPs and non-structural BMPs, and implementing LID components and techniques.

BMPs and LID components will be implemented by installing forebays and improving wetland areas at the East Inlet, redirecting storm water runoff into a proposed forebay and existing wetland at the South Inlet, installing maintenance access points and permanent dewatering ponds for collecting captured sediment at the East and South inlets, and implementing a winter sand reduction program within the watershed.

The City of Manchester Environmental Protection Department and Supplemental Environmental Project Plan Advisory Committee are providing the match required for this project.

**FIGURE INTERPRETATION**

- **Figure 1 and Table 1:** Figure 1 in Appendix A shows the historical and current year chlorophyll-a concentration in the water column. Table 1 in Appendix B lists the maximum, minimum, and mean concentration for each sampling year that the pond has been monitored through VLAP.

Chlorophyll-a, a pigment found in plants, is an indicator of the algal abundance. Because algae are usually microscopic plants that contain chlorophyll-a, and are naturally found in lake ecosystems, the chlorophyll-a concentration measured in the water gives an estimation of the algal concentration or lake productivity. **The median summer chlorophyll-a concentration for New Hampshire's lakes and ponds is 4.58 mg/m<sup>3</sup>.**

The current year data (the top graph) show that the chlorophyll-a concentration **decreased** from **June** to **July**, and then **increased** from **July** to **August**. The **elevated** chlorophyll concentration on the **June** and **August** sampling events indicated that an algal bloom had occurred.

The historical data (the bottom graph) show that the **2006** chlorophyll-a mean is **greater than** the state median and the similar lake median. For more information on the similar lake median, refer to Appendix F.

Overall, visual inspection of the historical data trend line (the bottom graph) shows a **variable** in-lake chlorophyll-a trend since monitoring began. Specifically the mean chlorophyll concentration has **fluctuated between approximately 10.81 and 24.84 mg/m<sup>3</sup>** since **2000**.

After 10 consecutive years of sample collection, we will be able to conduct a statistical analysis of the historical data to objectively determine if there has been a significant change in the annual mean chlorophyll-a concentration since monitoring began.

While algae are naturally present in all ponds, an excessive or increasing amount of any type is not welcomed. In freshwater ponds, phosphorus is the nutrient that algae typically depend upon for growth in New Hampshire lakes. Algal concentrations may increase as nonpoint sources of phosphorus from the watershed increase, or as in-lake phosphorus sources increase. Therefore, it is extremely important for volunteer monitors to continually educate all watershed residents about management practices that can be implemented to minimize phosphorus loading to surface waters.

- **Figures 2a and 2b and Tables 3a and 3b:** Figure 2a in Appendix A shows the historical and current year data for transparency without the use of a viewscope and Figure 2b shows the current year data for transparency with the use of a viewscope. Table 3a in Appendix B lists the maximum, minimum and mean transparency data without the use of a viewscope and Table 3b lists the maximum, minimum and mean transparency data with the use of a viewscope for each year that the pond has been monitored through VLAP.

Volunteer monitors use the Secchi disk, a 20 cm disk with alternating black and white quadrants, to measure how far a person can see into the water. Transparency, a measure of water clarity, can be affected by the amount of algae and sediment in the water, as well as the natural color of the water. **The median summer transparency for New Hampshire's lakes and ponds is 3.2 meters.**

The current year data (the top graph) show that the non-viewscope in-lake transparency **increased** from **June** to **July**, and then **decreased** from **July** to **August**.

It is important to note that as the chlorophyll concentration **decreased** from **June** to **July**, the transparency **increased**, and as the chlorophyll **increased** from **July** to **August**, the transparency **decreased**. We typically expect this **inverse** relationship in lakes. As the amount of algal cells in the water **decreases**, the depth to which one can see into the water column typically **increases**, and vice-versa.

The historical data (the bottom graph) show that the **2006** mean non-viewscope transparency is **less than** the state median and is **slightly less than** the similar lake median. Please refer to Appendix F for more information about the similar lake median.

The transparency measured with the viewscope was **equal to** or **greater than** the transparency measured without the viewscope on each sampling event this summer. As discussed previously, a comparison of the transparency readings taken with and without the use of a viewscope shows that the viewscope typically increases the depth to which the Secchi disk can be seen into the lake, particularly on sunny and windy days. We recommend that your group measure Secchi disk transparency with and without the viewscope on each sampling event.

It is important to note that viewscope transparency data is not compared to a New Hampshire median or similar lake median. This is because lake transparency has not been historically measured by DES with a viewscope. At some point in the future, the New

Hampshire and similar lake medians for viewscope transparency will be calculated and added to the appropriate graphs.

Overall, visual inspection of the historical data trend line (the bottom graph) shows a **variable** trend for in-lake non-viewscope transparency. Specifically, the transparency has **fluctuated between approximately 1.76 and 3.10 meters** since monitoring began in **2000**.

Again, please keep in mind that this trend is based on **seven** years of data. As previously discussed, after 10 consecutive years of sample collection, we will be able to conduct a statistical analysis of the historical data to objectively determine if there has been a significant change in the annual mean transparency since monitoring began.

Typically, high intensity rainfall causes sediment-laden stormwater runoff to flow into surface waters, thus increasing turbidity and decreasing clarity. Efforts should continually be made to stabilize stream banks, pond shorelines, disturbed soils within the watershed, and especially dirt roads located immediately adjacent to the edge of tributaries and the pond. Guides to best management practices that can be implemented to reduce, and possibly even eliminate, nonpoint source pollutants, are available from DES upon request.

- **Figure 3 and Table 8:** The graphs in Figure 3 in Appendix A show the amount of epilimnetic (upper layer) phosphorus and hypolimnetic (lower layer) phosphorus; the inset graphs show current year data. Table 8 in Appendix B lists the annual maximum, minimum, and median concentration for each deep spot layer and each tributary since the pond has been sampled through VLAP.

Phosphorus is typically the limiting nutrient for plant and algae growth in New Hampshire's lakes and ponds. Excessive phosphorus in a pond can lead to increased plant and algal growth over time.

**The median summer total phosphorus concentration in the epilimnion (upper layer) of New Hampshire's lakes and ponds is 12 ug/L. The median summer phosphorus concentration in the hypolimnion (lower layer) is 14 ug/L.**

The current year data for the epilimnion (the top inset graph) show that the phosphorus concentration **decreased** from **June to July**, and then **remained relatively stable** from **July to August**.

The **elevated** epilimnetic phosphorus concentration measured on the **June** sampling event may have been due to phosphorus-enriched stormwater runoff that flowed into the surface layer of the pond since

it rained approximately **two inches** during the **48 hours** prior to sampling.

The historical data show that the **2006** mean epilimnetic phosphorus concentration is **greater than** the state median and the similar lake median. Refer to Appendix F for more information about the similar lake median.

The current year data for the hypolimnion (the bottom inset graph) show that the phosphorus concentration **increased steadily** from **July** to **August**.

The turbidity of the hypolimnion (lower layer) sample was **elevated** on the **June** and **July** sampling events this year (**30 and 67.7 NTUs, respectively**). The hypolimnetic turbidity has been **at least slightly elevated** on many sampling events during previous years. This suggests that the pond bottom is covered by a thick organic layer of sediment which is easily disturbed. When the pond bottom is disturbed, sediment, which typically contains attached phosphorus, is released into the water column. When collecting the hypolimnion sample, make sure that there is no sediment in the Kemmerer Bottle before filling the sample bottles.

The historical data show that the **2006** mean hypolimnetic phosphorus concentration is **much greater than** the state median and the similar lake median. Please refer to Appendix F for more information about the similar lake median.

Overall, visual inspection of the historical data trend line for the epilimnion shows a **slightly variable** phosphorus trend. Specifically, the mean annual epilimnetic phosphorus concentration has **fluctuated between approximately 16.5 and 30.1 ug/L** since monitoring began in **2000**.

Overall, visual inspection of the historical data trend line for the hypolimnion shows a **highly variable** phosphorus trend since monitoring began. Specifically the mean annual concentration has **fluctuated between approximately 64 and 202 ug/L** since **2000**.

As discussed previously, after 10 consecutive years of sample collection, we will be able to conduct a statistical analysis of the historical data to objectively determine if there has been a significant change in the annual mean phosphorus concentration since monitoring began.

One of the most important approaches to reducing phosphorus loading to a waterbody is to continually educate watershed residents about the sources of phosphorus in a watershed and how excessive

phosphorus loading can negatively affect the ecology and the recreational, economical, and ecological value of lakes and ponds.

#### TABLE INTERPRETATION

##### ➤ **Table 2: Phytoplankton**

Table 2 in Appendix B lists the current and historical phytoplankton species observed in the pond. Specifically, this table lists the three most dominant phytoplankton species observed in the sample and their relative abundance in the sample.

The dominant phytoplankton species observed in the **June** sample were ***Asterionella* (diatom)**, ***Dinobryon* (golden-brown)**, and ***Synura* (golden-brown)**.

The dominant phytoplankton species observed in the **July** sample were ***Ceratium* (dinoflagellate)**, ***Staurastrum* (green)**, and ***Tabellaria* (diatom)**.

The dominant phytoplankton species observed in the **August** sample were ***Oscillatoria* (cyanobacteria)**, ***Ceratium* (dinoflagellate)**, and ***Fragilaria* (diatom)**.

Phytoplankton populations undergo a natural succession during the growing year. Please refer to the “Biological Monitoring Parameters” section of this report for a more detailed explanation regarding yearly plankton succession. Diatoms and golden-brown algae are typical in New Hampshire’s less productive lakes and ponds.

##### ➤ **Table 2: Cyanobacteria**

The cyanobacterium ***Oscillatoria*** was the **most-dominant** species observed in the **August** plankton sample. ***This species, if present in large amounts, can be toxic to livestock, wildlife, pets, and humans.*** Please refer to the “Biological Monitoring Parameters” section of this report for a more detailed explanation regarding cyanobacteria.

Cyanobacteria can reach nuisance levels when phosphorus loading from the watershed to surface waters is increased and favorable environmental conditions occur, such as a period of sunny, warm weather.

The presence of cyanobacteria serves as a reminder of the pond’s delicate balance. Watershed residents should continue to act proactively to reduce nutrient loading to the pond by eliminating

fertilizer use on lawns, keeping the pond shoreline natural, re-vegetating cleared areas within the watershed, and properly maintaining septic systems and roads.

In addition, residents should also observe the pond in September and October during the time of fall turnover (lake mixing) to document any algal blooms that may occur. Cyanobacteria have the ability to regulate their depth in the water column by producing or releasing gas from vesicles. However, occasionally lake mixing can affect their buoyancy and cause them to rise to the surface and bloom. Wind and currents tend to “pile” cyanobacteria into scums that accumulate in one section of the pond. If a fall bloom occurs, please collect a sample in any clean jar or bottle and contact the VLAP Coordinator.

➤ **Table 4: pH**

Table 4 in Appendix B presents the in-lake and tributary current year and historical pH data.

pH is measured on a logarithmic scale of 0 (acidic) to 14 (basic). pH is important to the survival and reproduction of fish and other aquatic life. A pH below 6.0 typically limits the growth and reproduction of fish. A pH between 6.0 and 7.0 is ideal for fish. The median pH value for the epilimnion (upper layer) in New Hampshire’s lakes and ponds is **6.6**, which indicates that the surface waters in the state are slightly acidic. For a more detailed explanation regarding pH, please refer to the “Chemical Monitoring Parameters” section of this report.

The mean pH at the deep spot this year ranged from **6.45** in the hypolimnion to **7.08** in the epilimnion, which means that the water is **slightly acidic** near the pond bottom and **approximately neutral** near the pond surface.

It is important to point out that the pH in the hypolimnion (lower layer) was **lower (more acidic)** than in the epilimnion (upper layer). This increase in acidity near the pond bottom is likely due to the decomposition of organic matter and the release of acidic by-products into the water column.

Due to the presence of granite bedrock in the state and acid deposition received from snowmelt, rainfall, and atmospheric particulates, there is not much that can be feasibly done to effectively increase pond pH.

➤ **Table 5: Acid Neutralizing Capacity**

Table 5 in Appendix B presents the current year and historical epilimnetic ANC for each year the pond has been monitored through VLAP.

Buffering capacity (ANC) describes the ability of a solution to resist changes in pH by neutralizing the acidic input. The median ANC value for New Hampshire's lakes and ponds is **4.9 mg/L**, which indicates that many lakes and ponds in the state are at least "moderately vulnerable" to acidic inputs. For a more detailed explanation about ANC, please refer to the "Chemical Monitoring Parameters" section of this report.

The mean acid neutralizing capacity (ANC) of the epilimnion (upper layer) was **19.3 mg/L**, which is ***much greater than*** the state median. In addition, this indicates that the pond has a ***low vulnerability*** to acidic inputs.

➤ **Table 6: Conductivity**

Table 6 in Appendix B presents the current and historical conductivity values for tributaries and in-lake data. Conductivity is the numerical expression of the ability of water to carry an electric current, which is determined by the number of negatively charged ions from metals, salts, and minerals in the water column. The median conductivity value for New Hampshire's lakes and ponds is **40.0 uMhos/cm**. For a more detailed explanation, please refer to the "Chemical Monitoring Parameters" section of this report.

The mean annual epilimnetic conductivity at the deep spot this year was **469.33 uMhos/cm**, which is ***much greater than*** the state median.

The conductivity continued to remain ***much greater than*** the state mean in the pond and inlets this season. Typically, elevated conductivity indicates the influence of pollutant sources associated with human activities. These sources include failed or marginally functioning septic systems, agricultural runoff, and road runoff which contains road salt during the spring snow-melt. New development in the watershed can alter runoff patterns and expose new soil and bedrock areas, which could also contribute to increasing conductivity. In addition, natural sources, such as iron and manganese deposits in bedrock, can influence conductivity.

We recommend that your monitoring group conduct a shoreline conductivity survey of the pond and the tributaries with ***elevated*** conductivity to help identify the sources of conductivity.

*To learn how to conduct a shoreline or tributary conductivity survey, please refer to the 2004 special topic article, which is posted on the VLAP website at [http://www.des.nh.gov/wmb/vlap/2004/documents/Appendix\\_D.pdf](http://www.des.nh.gov/wmb/vlap/2004/documents/Appendix_D.pdf) or contact the VLAP Coordinator.*

It is likely that de-icing materials applied to nearby roadways during the winter months may be influencing the conductivity in the pond. In New Hampshire, the most commonly used de-icing material is salt (sodium chloride).

*A limited amount of chloride sampling was conducted during **2005 and 2006**. Please refer to the discussion of **Table 13** for more information.*

➤ **Table 7a and Table 7b: Total Kjeldahl Nitrogen and Nitrite+Nitrate Nitrogen**

Table 7a in Appendix B presents the current year and historical Total Kjeldahl Nitrogen and Table 7b presents the current year and historical nitrite and nitrate nitrogen. Nitrogen is another nutrient that is essential for the growth of plants and algae. Nitrogen is typically the limiting nutrient in estuaries and coastal ecosystems. However, in freshwater, nitrogen is not typically the limiting nutrient. Therefore, nitrogen is not typically sampled through VLAP. However, if phosphorus concentrations in freshwater are elevated, then nitrogen loading may stimulate additional plant and algal growth. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

The pond is likely **phosphorus-limited**. Therefore, it is not critical to conduct nitrogen sampling.

➤ **Table 8: Total Phosphorus**

Table 8 in Appendix B presents the current year and historical total phosphorus data for in-lake and tributary stations. Phosphorus is the nutrient that limits the algae’s ability to grow and reproduce. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

The total phosphorus concentration in the **Inlet** was **elevated (160 ug/L)** on the **June** sampling event. The turbidity of the sample was **slightly elevated (2.09 NTUs)**. It had rained approximately **two inches** during the **48 hours** prior to sampling, which suggests that erosion is occurring in this area of the watershed.

We recommend that your monitoring group conduct a stream survey and rain event sampling along this tributary. This additional sampling may allow us to determine what is causing the **elevated** levels of turbidity and phosphorus.

*For a detailed explanation on how to conduct rain event sampling and stream surveys, please refer to the 2002 VLAP Annual Report special topic article, which is posted on the VLAP website at [http://www.des.nh.gov/wmb/vlap/2002/documents/Appndxd\\_monitoring.pdf](http://www.des.nh.gov/wmb/vlap/2002/documents/Appndxd_monitoring.pdf), or contact the VLAP Coordinator.*

- **Table 9 and Table 10: Dissolved Oxygen and Temperature Data**  
Table 9 in Appendix B shows the dissolved oxygen/temperature profile(s) collected during **2006**. Table 10 in Appendix B shows the historical and current year dissolved oxygen concentration in the hypolimnion (lower layer). The presence of dissolved oxygen is vital to fish and amphibians in the water column and also to bottom-dwelling organisms. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

The dissolved oxygen concentration was ***much lower in the hypolimnion (lower layer) than in the epilimnion (upper layer)*** at the deep spot on the **June, July, and August** sampling events. As ponds age, and as the summer progresses, oxygen typically becomes **depleted** in the hypolimnion by the process of decomposition. Specifically, the reduction of hypolimnetic oxygen is primarily a result of biological organisms using oxygen to break down organic matter, both in the water column and particularly at the bottom of the pond where the water meets the sediment. When hypolimnetic oxygen concentration is depleted to less than 1 mg/L, **as it was on each sampling event this year and on many previous annual visits**, the phosphorus that is normally bound up in the sediment may be re-released into the water column, a process referred to as ***internal phosphorus loading***.

Since an internal source of phosphorus in the pond may be present, it is even more important that watershed residents act proactively to minimize phosphorus loading from the watershed.

- **Table 11: Turbidity**  
Table 11 in Appendix B lists the current year and historical data for in-lake and tributary turbidity. Turbidity in the water is caused by suspended matter, such as clay, silt, and algae. Water clarity is strongly influenced by turbidity. Please refer to the “Other Monitoring Parameters” section of this report for a more detailed explanation.

As discussed previously, the hypolimnetic turbidity was **elevated** on the **June** and **July** sampling events this year, and has been **at least slightly elevated** on many sampling events during previous years. This suggests that the pond bottom is covered by a thick organic layer of sediment which is easily disturbed. When the pond bottom is disturbed, sediment, which typically contains attached phosphorus, is released into the water column. When collecting the hypolimnion sample, make sure that there is no sediment in the Kemmerer Bottle before filling the sample bottles.

Also discussed previously, the **epilimnetic** and **Inlet** turbidity on the **June** sampling event were **slightly elevated**, possibly due to sediment-laden stormwater runoff which flowed into the pond during a two inch rain event that occurring during the **48 hours** prior to sampling.

➤ **Table 12: Bacteria (*E.coli*)**

Table 12 in Appendix B lists the current year and historical data for bacteria (*E.coli*) testing. *E. coli* is a normal bacterium found in the large intestine of humans and other warm-blooded animals. *E.coli* is used as an indicator organism because it is easily cultured and its presence in the water, in defined amounts, indicates that sewage **may** be present. If sewage is present in the water, potentially harmful disease-causing organisms **may** also be present.

Bacteria sampling was not conducted this year. If residents are concerned about sources of bacteria such as failing septic systems, animal waste, or waterfowl waste, it is best to conduct *E. coli* testing when the water table is high, when beach use is heavy, or immediately after rain events.

➤ **Table 13: Chloride**

Table 13 in Appendix B lists the current year and the historical data for chloride sampling. The chloride ion (Cl<sup>-</sup>) is found naturally in some surfacewaters and groundwaters and in high concentrations in seawater. Research has shown that elevated chloride levels can be toxic to freshwater aquatic life. In order to protect freshwater aquatic life in New Hampshire, the state has adopted **acute and chronic** chloride criteria of **860 and 230 mg/L** respectively. The chloride content in New Hampshire lakes is naturally low, generally less than 2 mg/L in surface waters located in remote areas away from habitation. Higher values are generally associated with salted highways and, to a lesser extent, with septic inputs. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

Chloride sampling was **not** conducted during **2006**.

The deep spot was sampled for chloride during **2004** and **2005**. Many of the results **exceeded** the state chronic chloride criteria of 230 mg/L for surface waters, which suggests that deicing materials are leaching into the pond.

We recommend that your monitoring group conduct chloride sampling in the epilimnion at the deep spot and in the tributaries near salted roadways, particularly in the spring, soon during snow-melt and during rain events during the summer. This will establish a baseline of data that will assist your monitoring group and DES to determine lake quality trends in the future.

*Please note that there will be an additional cost for each of the chloride samples and that these samples must be analyzed at the DES laboratory in Concord.*

In addition, if your group is concerned about salt use on a particular roadway, we recommend contacting the town road agent or the Department of Transportation to discuss the implementation of a low-salt area near the pond and/or its major tributaries. We also recommend that your group work with watershed residents and business owners to reduce the application of chloride containing de-icing agents to driveways and walkways.

*To learn more about conductivity and chloride pollution and what can be done about to minimize it, please refer to the 2004 VLAP Annual Report special topic article, which is posted on the VLAP website at [http://www.des.state.nh.us/WMB/VLAP/2004/documents/Appendix\\_D.pdf](http://www.des.state.nh.us/WMB/VLAP/2004/documents/Appendix_D.pdf) or contact the VLAP Coordinator.*

- **Table 14: Current Year Biological and Chemical Raw Data**  
Table 14 in Appendix B lists the most current sampling year results. Since the maximum, minimum, and annual mean values for each parameter are not shown on this table, this table displays the current year “raw,” meaning unprocessed, data. The results are sorted by station, depth, and then parameter.
  
- **Table 15: Station Table**  
As of the spring of 2004, all historical and current year VLAP data are included in the DES Environmental Monitoring Database (EMD). To facilitate the transfer of VLAP data into the EMD, a new station identification system had to be developed. While volunteer

monitoring groups can still use the sampling station names that they have used in the past and are most familiar with, an EMD station name also exists for each VLAP sampling location. Table 15 in Appendix B identifies what EMD station name corresponds to the station names you have used in the past and will continue to use in the future.

#### **DATA QUALITY ASSURANCE AND CONTROL**

##### **Annual Assessment Audit:**

An annual DES biologist visit was not scheduled during 2006.

***Please contact the VLAP Coordinator during the spring of 2007 to schedule an annual biologist visit.***

##### **Sample Receipt Checklist:**

Each time your monitoring group dropped off samples at the laboratory this summer, the laboratory staff completed a sample receipt checklist to assess and document if your group followed proper sampling techniques when collecting the samples. The purpose of the sample receipt checklist is to minimize, and hopefully eliminate, future re-occurrences of improper sampling techniques.

Overall, the sample receipt checklist showed that your monitoring group did an **excellent** job when collecting samples and submitting them to the laboratory this year! Specifically, the members of your monitoring group followed the proper field sampling procedures and there was no need for the laboratory staff to contact your group with questions, and no samples were rejected for analysis.

#### **USEFUL RESOURCES**

*Acid Deposition Impacting New Hampshire's Ecosystems*, DES fact sheet ARD-32, (603) 271-2975 or [www.des.nh.gov/factsheets/ard/ard-32.htm](http://www.des.nh.gov/factsheets/ard/ard-32.htm).

*Best Management Practices to Control Nonpoint Source Pollution: A Guide for Citizens and Town Officials*, DES Booklet WD-03-42, (603) 271-2975.

*Best Management Practices for Well Drilling Operations*, DES fact sheet WD-WSEB-21-4, (603) 271-2975 or [www.des.nh.gov/factsheets/ws/ws-21-4.htm](http://www.des.nh.gov/factsheets/ws/ws-21-4.htm).

*Canada Geese Facts and Management Options*, DES fact sheet BB-53, (603) 271-2975 or [www.des.nh.gov/factsheets/bb/bb-53.htm](http://www.des.nh.gov/factsheets/bb/bb-53.htm).

*Cyanobacteria in New Hampshire Waters Potential Dangers of Blue-Green Algae Blooms*, DES fact sheet WMB-10, (603) 271-2975 or [www.des.nh.gov/factsheets/wmb/wmb-10.htm](http://www.des.nh.gov/factsheets/wmb/wmb-10.htm).

*Erosion Control for Construction in the Protected Shoreland Buffer Zone*, DES fact sheet WD-SP-1, (603) 271-2975 or [www.des.nh.gov/factsheets/sp/sp-1.htm](http://www.des.nh.gov/factsheets/sp/sp-1.htm).

*Impacts of Development Upon Stormwater Runoff*, DES fact sheet WD-WQE-7, (603) 271-2975 or [www.des.nh.gov/factsheets/wqe/wqe-7.htm](http://www.des.nh.gov/factsheets/wqe/wqe-7.htm).

*Lake Protection Tips: Some Do's and Don'ts for Maintaining Healthy Lakes*, DES fact sheet WD-BB-9, (603) 271-2975 or [www.des.nh.gov/factsheets/bb/bb-9.htm](http://www.des.nh.gov/factsheets/bb/bb-9.htm).

*Low Impact Development Hydrologic Analysis*. Manual prepared by Prince George's County, Maryland, Department of Environmental Resources. July 1999. To access this document, visit [www.epa.gov/owow/nps/lid\\_hydr.pdf](http://www.epa.gov/owow/nps/lid_hydr.pdf) or call the EPA Water Resource Center at (202) 566-1736.

*Low Impact Development: Taking Steps to Protect New Hampshire's Surface Waters*, DES fact sheet WD-WMB-16, (603) 271-2975 or [www.des.nh.gov/factsheets/wmb/wmb-17.htm](http://www.des.nh.gov/factsheets/wmb/wmb-17.htm).

*Proper Lawn Care In the Protected Shoreland, The Comprehensive Shoreland Protection Act*, DES fact sheet WD-SP-2, (603) 271-2975 or [www.des.nh.gov/factsheets/sp/sp-2.htm](http://www.des.nh.gov/factsheets/sp/sp-2.htm).

*Road Salt and Water Quality*, DES fact sheet WD-WMB-4, (603) 271-2975 or [www.des.nh.gov/factsheets/wmb/wmb-4.htm](http://www.des.nh.gov/factsheets/wmb/wmb-4.htm).

*Sand Dumping - Beach Construction*, DES fact sheet WD-BB-15, (603) 271-2975 or [www.des.nh.gov/factsheets/bb/bb-15.htm](http://www.des.nh.gov/factsheets/bb/bb-15.htm).



*Shorelands Under the Jurisdiction of the Comprehensive Shoreland Protection Act*, DES fact sheet SP-4, (603) 271-2975 or [www.des.nh.gov/factsheets/sp/sp-4.htm](http://www.des.nh.gov/factsheets/sp/sp-4.htm).

*Soil Erosion and Sediment Control on Construction Sites*, DES fact sheet WQE-6, (603) 271-2975 or [www.des.nh.gov/factsheets/wqe/wqe-6.htm](http://www.des.nh.gov/factsheets/wqe/wqe-6.htm).

*Watershed Districts and Ordinances*, DES fact sheet WD-WMB-16, (603) 271-2975 or [www.des.nh.gov/factsheets/wmb/wmb-16.htm](http://www.des.nh.gov/factsheets/wmb/wmb-16.htm).



# Nutt Pond, Manchester

**Figure 1.** Monthly and Historical Chlorophyll-a Results




# Nutt Pond, Manchester

**Figure 2a.** Monthly and Historical Transparency without Viewscope Results




# Nutt Pond, Manchester

**Figure 2b.** Monthly and Historical Transparency with Viewscope Results



2006 Transparency with Viewscope Results



Historical Transparency with Viewscope Results

# Nutt Pond, Manchester

**Figure 3.** Monthly and Historical Total Phosphorus Data.

